

## **Authors**

Slawa Rokicki Rutgers School of Public Health

Jessica Cohen Harvard University

Joshua A. Salomon Harvard School of Public Health

Günther Fink Swiss Tropical and Public Health Institute

## Impact of a Text Messaging Program on Adolescent Reproductive Health: A Cluster-Randomized Trial in Ghana\*

Slawa Rokicki<sup>2</sup>, PhD, Jessica Cohen<sup>2</sup>, PhD, Joshua A, Salomon<sup>2</sup>, PhD, and Günther Fink<sup>2</sup>, PhD

## Abstract

Objectives. To evaluate whether text-messaging programs can improve reproductive health among adolescent girls in low- and middle-income countries.

Methods. We conducted a cluster-randomized controlled trial among 756 female students aged 14 to 24 years in Accra, Ghana, in 2014. We randomized 38 schools to unidirectional intervention (n=12), interactive intervention (n=12), and control (n=14). The unidirectional intervention sent participants text messages with reproductive health information. The interactive intervention engaged adolescents in text-messaging reproductive health quiz games. The primary study outcome was reproductive health knowledge at 3 and 15 months. Additional outcomes included self-reported pregnancy and sexual behavior. Analysis was by intent-to-treat.

Results. From baseline to 3 months, the unidirectional intervention increased knowledge by 11 percentage points (95% confidence interval [CI]=7, 15) and the interactive intervention by 24 percentage points (95% CI=19, 28), from a control baseline of 26%. Although we found no changes in reproductive health outcomes overall, both unidirectional (odds ratio 10R1=0.14: 959

## Impact of a Text Messaging Program on Adolescent Reproductive Health: A Cluster-Randomized Trial in Ghana

*Objectives*. To evaluate whether text-messaging programs can improve reproductive health among adolescent girls in low- and middle-income countries.

Methods. We conducted a cluster-randomized controlled trial among 756 female students

<sup>\*</sup> Published online at: http://ajph.aphapublications.org/doi/abs/10.2105/AJPH.2016.303562

<sup>&</sup>lt;sup>3</sup> Interfaculty Initiative in Health Policy, Harvard University Geary Institute for Public Policy, University College Dublin. Email: slawa.rokicki@cod.ie

<sup>&</sup>lt;sup>2</sup> Department of Global Health and Population, Harvard T.H. Chun School of Public Health



aged 14 to 24 years in Accra, Ghana, in 2014. We randomized 38 schools to unidirectional intervention (n=12), interactive intervention (n=12), and control (n=14). The unidirectional intervention sent participants text messages with reproductive health information. The interactive intervention engaged adolescents in text-messaging reproductive health quizzes. The primary study outcome was reproductive health knowledge at 3 and 15 months. Additional outcomes included self-reported pregnancy and sexual behavior. Analysis was by intent-to-treat.

Results. From baseline to 3 months, the unidirectional intervention increased knowledge by 11 percentage points (95% confidence interval [CI] = 7, 15) and the interactive intervention by 24 percentage points (95% CI = 19, 28), from a control baseline of 26%. Although we found no changes in reproductive health outcomes overall, both unidirectional (odds ratio [OR] = 0.14; 95% CI = 0.03, 0.71) and interactive interventions (OR = 0.15; 95% CI = 0.03, 0.86) lowered odds of self-reported pregnancy for sexually active participants.

December 20, 2016