

Authors

Slawa Rokicki
Rutgers School of Public Health

Jessica Cohen
Harvard University

Joshua A. Salomon
Harvard School of Public Health

Günther Fink
Swiss Tropical and Public Health Institute

Impact of a Text Messaging Program on Adolescent Reproductive Health: A Cluster-Randomized Trial in Ghana*

Slawa Rokicki¹, PhD, Jessica Cohen², PhD, Joshua A. Salomon², PhD, and Günther Fink², PhD

Abstract

Objectives. To evaluate whether text-messaging programs can improve reproductive health among adolescent girls in low- and middle-income countries.

Methods. We conducted a cluster-randomized controlled trial among 756 female students aged 14 to 24 years in Accra, Ghana, in 2014. We randomized 38 schools to unidirectional intervention (n=12), interactive intervention (n=12), and control (n=14). The unidirectional intervention sent participants text messages with reproductive health information. The interactive intervention engaged adolescents in text-messaging reproductive health quiz games. The primary study outcome was reproductive health knowledge at 3 and 15 months. Additional outcomes included self-reported pregnancy and sexual behavior. Analysis was by intent-to-treat.

Results. From baseline to 3 months, the unidirectional intervention increased knowledge by 11 percentage points (95% confidence interval [CI]=7, 15) and the interactive intervention by 24 percentage points (95% CI=19, 28), from a control baseline of 26%. Although we found no changes in reproductive health outcomes overall, both unidirectional (odds ratio [OR]=0.14; 95%

* Published online at: <http://ajph.aphapublications.org/doi/abs/10.2105/AJPH.2016.303562>

¹ Interfaculty Initiative in Health Policy, Harvard University
Geary Institute for Public Policy, University College Dublin. Email: slawa.rokicki@ucd.ie

² Department of Global Health and Population, Harvard T.H. Chan School of Public Health

Impact of a Text Messaging Program on Adolescent Reproductive Health: A Cluster-Randomized Trial in Ghana

Objectives. To evaluate whether text-messaging programs can improve reproductive health among adolescent girls in low- and middle-income countries.

Methods. We conducted a cluster-randomized controlled trial among 756 female students

aged 14 to 24 years in Accra, Ghana, in 2014. We randomized 38 schools to unidirectional intervention ($n = 12$), interactive intervention ($n = 12$), and control ($n = 14$). The unidirectional intervention sent participants text messages with reproductive health information. The interactive intervention engaged adolescents in text-messaging reproductive health quizzes. The primary study outcome was reproductive health knowledge at 3 and 15 months. Additional outcomes included self-reported pregnancy and sexual behavior. Analysis was by intent-to-treat.

Results. From baseline to 3 months, the unidirectional intervention increased knowledge by 11 percentage points (95% confidence interval [CI] = 7, 15) and the interactive intervention by 24 percentage points (95% CI = 19, 28), from a control baseline of 26%. Although we found no changes in reproductive health outcomes overall, both unidirectional (odds ratio [OR] = 0.14; 95% CI = 0.03, 0.71) and interactive interventions (OR = 0.15; 95% CI = 0.03, 0.86) lowered odds of self-reported pregnancy for sexually active participants.

December 20, 2016