

Authors

Günther Fink
Swiss Tropical and Public Health Institute

Rachel Levenson
Harvard University

Peter Rockers
Boston University

HEALTH | POLICY BRIEF

Poster Child for Healthy Growth

In rural Zambia, simple, low-cost growth charts reduced malnutrition.



JULY 2018



Key Findings

- Simple growth charts, which allowed parents to see if their child had a normal height for their age, did not reduce growth deficits on average among the 547 children in the study, but among malnourished children, reduced stunting by 22 percentage points.
- In contrast, inviting caregivers to quarterly meetings to learn if their children had a normal height and weight and providing food supplements to malnourished children had no impact on rates of stunting.
- Neither home-based growth charts nor community-based monitoring were found to impact children's cognitive development.
- Home-based growth charts appear to be a cost-effective tool to reduce physical growth deficits in this context. For every dollar that was invested in growth charts, children who otherwise would have been stunted gained an estimated \$16 in additional lifetime wages.

RESEARCHERS

Günther Fink (Swiss Tropical and Public Health Institute)
Rachel Levenson (Innovations for Poverty Action)
Sarah Tembo (Innovations for Poverty Action)
Peter Rockers (Boston University)

FUNDERS

The Bill & Melinda Gates Foundation

SOURCE

Fink, Günther, Rachel Levenson, Sarah Tembo, and Peter C. Rockers. "Home and community-based growth monitoring to reduce early-life growth faltering among children: randomised controlled trial." *The American journal of clinical nutrition* (2017): 491-97945.

COUNTRY

Zambia

TIME PERIOD

2014-2015

SAMPLE

127 villages (547 children 6-24 months old)

Poster Child for Healthy Growth

Key Findings:

- Simple growth charts, which allowed parents to see if their child had a normal height for their age, did not reduce reduce growth deficits on average among the 547 children in the study, but among malnourished children, reduced stunting by 22 percentage points.
- In contrast, inviting caregivers to quarterly meetings to learn if their children had a normal height and weight and providing food supplements to malnourished children had no impact on rates of stunting.
- Neither home-based growth charts nor community-based monitoring were found to impact children's cognitive development.

- Home-based growth charts appear to be a cost-effective tool to reduce physical growth deficits in this context. For every dollar that was invested in growth charts, children who otherwise would have been stunted gained an estimated \$16 in additional lifetime wages.

July 10, 2018